

Equilibrium Operational Space and Stability Limits in Early Operation of MAST-U

J.W. Berkery (PPPL)





S.A. Sabbagh, V. Zamkovska, J. Butt (Columbia University)

C. Ham, L. Kogan, S. Gibson, D. Ryan, J. Harrison, S. Henderson, B. Patel, R. Scannell, R. Sharma, E. Trier, T. Wilson, and the MAST-U Team (UKAEA)

**Presented at the 21st International Spherical Torus Workshop,
November 2, 2022**

Supported by U.S. Department of Energy grants DE-SC0018623 and DE-AC02-09CH11466

- 750kA plasmas with 1s flat-top
- Good pedestal performance
- Conventional and Super-X divertors



3D VALEN model of surfaces to calibrate







2D wall current model



Magnetic reconstructions



Plus motional Stark effect



- Kinetic reconstructions forthcoming

[J.W. Berkery, *et al.*, PPCF **63**, 055014 (2021)]


[L. Kogan, *et al.*, EPS (2022)]

- Database includes 855 discharges (June-Oct, 2021)
 - Not every discharge – need good flattop, good reconstruction
 - Flattop split in 5ms intervals – 108,826 total equilibria
- MAST-U discharges were categorized in discrete allowed scenarios
 - Plasma current level: 400-450 kA, 600 kA, 700-750 kA
 - Neutral beam heating: Ohmic (OH), one beam (SW or SS), two beam (2B)
 - Divertor shape: conventional (CD), SuperX (SX), or unlabelled (UN)

I_p (kA)	Divertor	Ohmic	SW	SS	2B
400	CD	148		12	
	SX	64			
	UN	44			
600	CD	39	9	25	7
	SX	56	12		8
	UN	16		6	
700	CD	53	77	31	108
	SX	68			
	UN	30	31	3	9

Hugill diagram shows that MAST-U has yet to access low density, low q disruptive space




5



- Disruption Event Characterization and Forecasting (DECAF) code used
 - Produces disruptivity or operational space plots – here 50x50 grids
- Hugill diagram showed high disruptivity for MAST at low density, low q_{95}
 - MAST-U has yet to really access this region of operational space much


Before operation, MAST-U projected equilibria pressure and q_0 were scanned to find global stability limits

6



- Not all these p and q profile combinations are realizable in MASTU
- They serve to explore the parameter space


[J.W. Berkery, *et al.*, PPCF **62**, 085007 (2020)]




- Contours of $-\delta W$: red above the no-wall limit and blue below
- In each case, $\beta_N^{\text{no-wall}}$ increases slightly with internal inductance
 - $\beta_N^{\text{no-wall}} \sim 4$

[J.W. Berkery, *et al.*, PPCF **62**, 085007 (2020)]

MAST-U operational space



MAST-U operational space




Contours
contain ≥ 30
equilibria

Two beam
One beam, on axis
One beam, off axis
Ohmic

- Not surprisingly, two beam injection gives the highest β_N
 - The SS beam is oriented to deposit energy on-axis, so can generally lead to higher β_N plasmas than the off-axis SW beam
- Projected MAST-U no-wall limit: $\beta_N \sim 4$ and $\beta_N / I_i \sim 7$


[J.W. Berkery, *et al.*, PPCF **62**, 085007 (2020)]



- With the exception of vertical stability, increased elongation in STs tends to increase the stability of the plasmas
 - κ increases both β_t for fusion gain and $f_{bs} \sim \beta_p$ for pulse length, sustainment
- Increased elongation is a focus of the second campaign

MAST-U has generally stayed well under the Greenwald limit in the flattop to this point


10



- There were occasional cases of flattop plasmas approaching the Greenwald limit
 - Experiments in the second campaign will push to establish density limits
- Lower plasma current shots need to maintain lower density

In MAST-U plasmas the Greenwald limit can be avoided by quickly shrinking the plasma


11



- MAST rampdown Greenwald density limits happened often
 - Reducing I_p brought the limit down while the density was still rising
- In some MAST-U cases, the limit rises in the rampdown
 - The plasma's minor radius is rapidly decreasing

Edge density limits can be reached before the Greenwald limit, potentially giving an early warning

12



- GWL low, but edge high -> disruption
- Not stringent criteria
- In DECAF code could be one of multiple warnings

Giacomin Bernert

$$n_{\text{lim}} = \alpha A^{1/6} a^{3/14} P_{\text{SOL}}^{10/21} R_0^{-43/42} q^{-22/21} (1 + \kappa^2)^{-1/3} B_T^{2/3}$$

$$\bar{n}_{e,H5,\text{reg.}} = 0.506 \pm 0.192 \frac{P_{\text{heat}}^{0.396 \pm 0.13} I_p^{0.265 \pm 0.14}}{q_{95}^{0.323 \pm 0.14}},$$

- Various plasma scenarios achieved:
 - Ohmic, one, or two neutral beams, 400-750kA, CD or SX
- Magnetics-only equilibria produced and used by DECAF for operational space diagrams
- Generally, MAST-U still has margin to increase performance without encountering stability limits
 - Beta, towards projected global stability limits
 - Elongation, towards vertical stability limits
 - Density, towards Greenwald limit