

# Electron heating during neutral beam injection at Globus-M2.

Zhiltsov N.S.<sup>1</sup>, Kurskiev G.S.<sup>1</sup>, Bakharev N.N.<sup>1</sup>, Balachenkov I.M.<sup>1</sup>, Filippov S.V.<sup>1</sup>, Gusev V.K.<sup>1</sup>,  
Khromov N.A.<sup>1</sup>, Kiselev E.O.<sup>1</sup>, Koval A.N.<sup>1</sup>, Krikunov S.V.<sup>1</sup>, Minaev V.B.<sup>1</sup>, Miroshnikov I.V.<sup>1</sup>,  
Mukhin E.E.<sup>1</sup>, Novokhatsky A.N.<sup>1</sup>, Patrov M.I.<sup>1</sup>, Petrov Yu.V.<sup>1</sup>, Sakharov N.V.<sup>1</sup>, Shchegolev P.B.<sup>1</sup>,  
Shulyatiev K.D.<sup>1</sup>, Skrekel O.M.<sup>1</sup>, Solokha V.V.<sup>1</sup>, Solovey V.A.<sup>2</sup>, Telnova A.Yu.<sup>1</sup>, Tkachenko E.E.<sup>1</sup>,  
Tokarev V.A.<sup>1</sup>, Tolstyakov S.Yu.<sup>1</sup>, Tukhmeneva E.A.<sup>1</sup>, Varfolomeev V.I.<sup>1</sup>, Voronin A.V.<sup>1</sup>,  
Goryainov V.Yu.<sup>1</sup>, Yashin A.Yu.<sup>1, 3</sup>

<sup>1</sup> *Ioffe Institute, Saint-Petersburg, Russia*

<sup>2</sup> *Petersburg Nuclear Physics Institute named by B.P. Konstantinov of NRC, “Kurchatov Institute”, Saint-Petersburg, Russia*


<sup>3</sup> *Peter the Great St. Petersburg Polytechnic University, St. Petersburg, Russia*

E-mail: [nisovru@gmail.com](mailto:nisovru@gmail.com)



## Globus-M2 equatorial cross-section

| Globus-M2<br>Spherical tokamak |
|--------------------------------|
| $B_T = 0.9 \text{ T}$          |
| $I_p = 0.4 \text{ MA}$         |
| $R = 0.36 \text{ m}$           |
| $a = 0.24 \text{ m}$           |
| $R/a = 1.5$                    |
| Graphite wall                  |





## Globus-M2 equatorial cross-section

### Globus-M2 Spherical tokamak

$$B_T = 0.9 \text{ T}$$

$$I_p = 0.4 \text{ MA}$$

$$R = 0.36 \text{ m}$$

$$a = 0.24 \text{ m}$$


$$R/a = 1.5$$

**Graphite wall**

**NBI<sub>E=45keV</sub>**

$$P = 0.75 \text{ MW}$$

$$E = 45 \text{ keV}$$






# NBI<sub>E=45keV</sub> experiment setup

## Globus-M2 equatorial cross-section


| Globus-M2<br>Spherical tokamak |
|--------------------------------|
| $B_T = 0.9 \text{ T}$          |
| $I_p = 0.4 \text{ MA}$         |
| $R = 0.36 \text{ m}$           |
| $a = 0.24 \text{ m}$           |
| $R/a = 1.5$                    |
| Graphite wall                  |
| NBI <sub>2</sub>               |
| $P = 0.75 \text{ MW}$          |
| $E = 45 \text{ keV}$           |






# NBI<sub>E=45keV</sub> experiments

- NBI started at  $I_p$  ramp-up
- $n_e$  varied in range  $0.7 \cdot 10^{19} - 7.5 \cdot 10^{19} \text{ (m}^{-3}\text{)}$
- $T_e$  is independent from  $\langle n_e \rangle$  except for low  $\langle n_e \rangle$  (red).
- $W_e$  proportional to  $\langle n_e \rangle$ , absence of saturation







- peaked  $n_e$  profiles
- $n_e$  gradient increase with  $\langle n_e \rangle$  for  $r/a \leq 0.9$
- $n_e$  edge pedestal is rises for high  $\langle n_e \rangle$



# NBI<sub>E=45keV</sub> experiments




- peaked  $n_e$  profiles
- $n_e$  gradient increase with  $\langle n_e \rangle$  for  $r/a \leq 0.9$
- $n_e$  edge pedestal is rises for high  $\langle n_e \rangle$



- $T_e$  gradient is independent from  $\langle n_e \rangle$
- Central  $T_e = 1.3$  (keV) at  $n_e = 1 \cdot 10^{20} \text{ (m}^{-3}\text{)}$




# NBI<sub>E=45keV</sub> experiments




- Linear  $W_e$  ( $\langle n_e \rangle$ ) rise in NBI<sub>E=45</sub> regime
- NBI<sub>E=45keV</sub> regime has 175% of ohmic  $W_e$  at high  $\langle n_e \rangle$
- **Electron heating is evident for NBI<sub>E=45keV</sub> regime**




# NBI<sub>E=45keV</sub> experiments




- Linear  $W_e$  ( $\langle n_e \rangle$ ) rise in NBI<sub>E=45</sub> regime
- NBI<sub>E=45keV</sub> regime has 175% of ohmic  $W_e$  at high  $\langle n_e \rangle$
- **Electron heating is evident for NBI<sub>E=45keV</sub> regime**





- Diamagnetic measurements allow to estimate  $W_i$   
as  $W_{DIA} = W_e + W_i + W_{fast \perp}$



- Diamagnetic measurements allow to estimate  $W_i$   
as  $W_{\text{DIA}} = W_e + W_i + W_{\text{fast } \perp}$




- Diamagnetic measurements allow to estimate  $W_i$   
as  $W_{DIA} = W_e + W_i + W_{fast \perp}$
- anomalous  $\chi_i$



- Diamagnetic measurements allow to estimate  $W_i$   
as  $W_{DIA} = W_e + W_i + W_{fast \perp}$
- anomalous  $\chi_i$

- NBI<sub>E=45keV</sub> experiments point at low  $T_i$  values




$\tau_E = 11(\text{ms})$  for  $\langle n_e \rangle \sim 7 \cdot 10^{19} \text{ (m}^{-3}\text{)}$  and NBI<sub>E=45keV</sub> fits previous results for NBI<sub>E=28keV</sub>

Anomalous  $\chi_i$  leads to saturation of  $\tau_E$



# NBI<sub>1</sub> + NBI<sub>2</sub> experiment setup


## Globus-M2 equatorial cross-section





# NBI<sub>1</sub> + NBI<sub>2</sub> experiments

Globus-M2  $B_T = 0.9$  T,  $I_p = 0.35$ - $0.4$  MA, NBI<sub>1</sub> + NBI<sub>2</sub>





Hot ion mode achieved in wide  $\langle n_e \rangle$  range 0.15 - 1.6 ( $10^{20} \cdot \text{m}^{-3}$ )

$T_i = 4$  (keV) at  $\langle n_e \rangle = 4.9 \cdot 10^{19} \text{ (m}^{-3}\text{)}$



# NBI<sub>1</sub> + NBI<sub>2</sub> experiments

Globus-M2  $B_T = 0.9$  T,  $I_p = 0.35$ - $0.4$  MA, NBI<sub>1</sub> + NBI<sub>2</sub>




Hot ion mode achieved in wide  $\langle n_e \rangle$  range  $1.5 \cdot 10^{19}$  -  $1.6 \cdot 10^{20} \text{ (m}^{-3})$

$T_i = 4 \text{ (keV)}$  at  $\langle n_e \rangle = 4.9 \cdot 10^{19} \text{ (m}^{-3})$

$T_i / T_e$  up to 3



# NBI<sub>1</sub> + NBI<sub>2</sub> experiments





If we roughly top estimate  $W_i$  as  $W_e \cdot T_i / T_e$ , then:

- $W_i$  saturates for  $\langle n_e \rangle \geq 0.5 \cdot 10^{20}$  (m<sup>-3</sup>)
- Further  $W_p$  rise is due to  $W_e$



# NBI<sub>1</sub> + NBI<sub>2</sub> experiments




If we roughly top estimate  $W_i$  as  $W_e \cdot T_i / T_e$ , then:

- $W_i$  saturates for  $\langle n_e \rangle \geq 0.5 \cdot 10^{20} \text{ (m}^{-3}\text{)}$
- Further  $W_p$  rise is due to  $W_e$




# NBI<sub>1</sub> + NBI<sub>2</sub> experiments



If we roughly top estimate  $W_i$  as  $W_e \cdot T_i/T_e$ , then:

- $W_i$  saturates for  $\langle n_e \rangle \geq 0.5 \cdot 10^{20}$  (m<sup>-3</sup>)
- Further  $W_p$  rise is due to  $W_e$



- Same linear dependency as for NBI<sub>E=45keV</sub> mode at low  $\langle n_e \rangle$
- $\langle n_e \rangle = 5 \cdot 10^{19}$  (m<sup>-3</sup>)  $\Rightarrow$  slope changes
- $T_e > 1.3$  (keV) for  $\langle n_e \rangle = 1.7 \cdot 10^{20}$  (m<sup>-3</sup>)
- $W_e$  is independent from I<sub>p</sub> high  $\langle n_e \rangle$



# NBI<sub>1</sub> + NBI<sub>2</sub> experiments

Double NBI scenario:

NBI<sub>2</sub> from 0.125 (s) to 0.23 (s)

$I_p$  flattop from 0.175 (s) to 0.24 (s)

NBI<sub>1</sub> from 0.19 (s) to 0.22 (s)

L-H transition at 0.17 (s)

$D_\alpha$  intensity drop

type 5 ELMs


$\langle n_e \rangle$  from  $3 \cdot 10^{19} \text{ (m}^{-3}\text{)}$  to  $5.5 \cdot 10^{19} \text{ (m}^{-3}\text{)}$  on  $I_p$  flattop

$T_i$  measurements on NBI<sub>1</sub> beam by CXRS and NPA matches

$T_i(0)$  rises up to 4 (keV)


$T_e(0) = 1.6 \text{ (keV)}$

$W_{dia}$  matches  $W_e^{TS} + W_i^{CXRS} + W_{fast \perp}$






# NBI<sub>1</sub> + NBI<sub>2</sub> experiments




- $T_i(0) = 4$  (keV)
- $T_i(0) / T_e(0) = 2.7$
- Hot ion mode achieved with:  $B_T = 0.9$  (T)  
 $I_p = 0.35$  (MA)  
 $P_{\text{NBI } 1+2} = 0.7$  (MW)




# NBI<sub>1</sub> + NBI<sub>2</sub> experiments



- $T_i(0) = 4$  (keV)
- $T_i(0) / T_e(0) = 2.7$
- Hot ion mode achieved with:  $B_T = 0.9$  (T)  
 $I_p = 0.35$  (MA)  
 $P_{\text{NBI } 1+2} = 0.7$  (MW)



- $n_e(0) = 7.8 \cdot 10^{19}$  (m<sup>-3</sup>)
  - Peaked  $n_e$  profile
- Legend:
- NBI particle source
  - Improved central confinement



- Suppressed anomalous  $\chi_i$ .  
 $\chi_i \leq 1 \text{ (m}^2 \text{s}^{-1}\text{)}$  is close to neoclassical

- $\tau_E = 14.3 \text{ (ms)}$

- Typical for Globus-M2  $\chi_e \sim 2$  value

- $n_e(0) \cdot T_i(0) \cdot \tau_E = 4.3 \cdot 10^{18} \text{ (m}^{-3} \text{ keV s)}$



## Globus-M2, NBI<sub>2</sub> experiment:

- NBI<sub>E=45keV</sub> is heating electrons well  
 $T_e = 1.3$  (keV) in NBI<sub>E=45keV</sub> regime is twice as in ohmic regime for  $\langle n_e \rangle = 7.3 \cdot 10^{20} \text{ (m}^{-3}\text{)}$
- Cold ions indicated by joint analysis of diamagnetic and TS data

## Globus-M2, NBI<sub>1</sub>+ NBI<sub>2</sub> experiment:

Achieved:

- Hot ion mode for  $\langle n_e \rangle = [0.15 - 1.6] (10^{20} \cdot \text{m}^{-3})$
- $T_i = 4$  (keV)
- $T_i / T_e$  up to 3
- $n_e(0) \cdot \tau_E = 1.1 \cdot 10^{18} \text{ (m}^{-3} \text{ s)}$