


Linear Gyrokinetic Stability Analysis of ST40 Hot Ion Plasmas

Y. Ren¹, S. Kaye¹, M.S. Anastopoulos Tzanis², M. Sertoli², A. Dnestrovskii², M. Romanelli²

¹ Princeton Plasma Physics Laboratory, Princeton, NJ 08543
² Tokamak Energy Ltd, 173 Brook Drive, Milton Park, Oxfordshire OX14 4SD




Introduction

- ST40 is the first high-field spherical tokamak, achieving $I_p \approx 0.4\text{--}0.8$ MA, on-axis $B_T \approx 1.5\text{--}2.2$ T, and pulse durations ≈ 0.2 s in the most recent operations.
 - 2 MW of neutral beam heating is available: one source at 25 keV for 0.1 second; the other at 55 keV for 1 second
- The high ion temperature ST40 plasmas ($T_i \gg T_e \sim 3$ keV) were achieved, e.g., with $R=0.45$ m, $\kappa = 1.5$, aspect ratio $A=1.65$, $I_p = 0.6$ MA, $B_T \approx 1.9$ T, and a typical pulse length of 0.1 s.
 - Plasma profiles inferred using integrated analysis workflow that incorporated the line-of-sight integrated and profile measurements of the electron, ion temperatures and toroidal rotation, the line-of-sight integrated density measurements, and equilibrium constraints.
- It is imperative to explore operating micro-instabilities in ST40 plasmas, particularly in the fusion-relevant hot-ion regime.
 - Micro-turbulence considered to be a major candidate in driving anomalous transport in fusion plasmas, and the energy confinement time of ST40 most likely being limited by micro-turbulence.

Linear Stability is Studied at Multiple Radii in a ST40 Hot-ion Shot 9831

- $B_T \sim 1.8$ T, $I_p \sim 500$ kA and not highly shaped with $\kappa \sim 1.3$ and $\delta \sim 0.2$
- Impurity temperature, T_{im} , ~ 10 keV in the core, and ion temperature, T_i , is 1-1.5 keV lower than T_{im} in plasma center, clearly in the hot-ion regime
 - $T_{im} = T_i$ should be well satisfied in the outer half of the plasma.
- $\eta_e = \frac{L_{ne}}{L_{Te}}$ decreases from -1 to -0.5 from $\rho=0.8$ to 0.3, showing stabilization of ETG mode towards the plasm core.
- $\eta_i = \frac{L_{ni}}{L_{Ti}}$ increases from -1 to -1.5 from $\rho=0.8$ to 0.3, indicating ion mode more important towards the plasma core.
- The analysis was carried out at $\rho=0.3, 0.6, 0.8$ (ρ is the square root of normalized toroidal flux)



Significant Stabilization of Micro-instabilities towards Plasma Core


- Linear growth rates of both ion- and electron-scale ($k_\theta \rho_s \gtrsim 0.2$) modes decrease from the edge towards the core of the plasma
- Complete suppression of electron-scale modes ($k_\theta \rho_s \gtrsim 5$) at $\rho = 0.3$; No-monotonic for $k_\theta \rho_s < 0.2$
- ExB shearing rate $<$ the maximum γ in the ion scale, implying the importance of ion-scale modes in driving transport
- Ion-scale modes are mainly in the ion direction and electron scale modes are mainly in the electron direction
 - At $\rho=0.8$, there is a change in the propagation direction at $k_\theta \rho_s \approx 0.34$
- What are these ion-scale modes?
- Small- ω k ranges in the intermediate scale are seen from $k_\theta \rho_s \approx 5$ to 60 at $\rho = 0.8$ and from $k_\theta \rho_s \approx 5$ to 30 at $\rho = 0.6$ with ω only weakly increasing in amplitude with k
- Typical electrostatic ETG mode dispersion is only seen at higher wavenumbers, $\gtrsim \rho_e^{-1}$ with ω strongly increasing in amplitude with k
- What are these small- ω electron-scale modes?

Ubiquitous Mode (UM) is due to Nonadiabatic Responses of Both Trapped Electrons and Passing Ions


- This branch of solution of the dispersion equation is of TEM at lower k and is of UM at higher k with a purely growing fluid-like mode in the transition
 - Both TEM and ubiquitous mode are driven by wave-particle resonances
 - The fluid-like mode is non-resonantly driven by magnetic drifts of both trapped electrons and passing ions in the bad curvature region
- Nonadiabatic ion response in the electron scale is important
 - Adiabatic ions are usually assumed for electron scale, i.e. for the well-known ETG mode
 - Adiabatic ion response arises from $\frac{1}{k_\perp} \frac{\partial}{\partial b}$ term in the nonadiabatic ion response where $\frac{1}{k_\perp} \frac{\partial}{\partial b} \sim \frac{1}{\sqrt{b}}$ with large b and $b = \frac{k_\perp^2 \rho_s^2}{2}$
 - However, in local approximation, the non-adiabatic part of ion response is:

$$\frac{\omega + \omega_{di}}{\omega - \omega_{di}} \frac{1}{k_\perp} \frac{\partial}{\partial b} \frac{e \phi}{T_i n_0}$$
 where ω_{di} is the ion diamagnetic drift frequency and ω_{di} is the ion magnetic drift frequency.
 - If $\omega_{di} \ll \omega \ll \omega_{ci}$, with large b, the non-adiabatic part of ion response is written as


$$\frac{\omega_{ci}}{\omega} \frac{1}{k_\perp} \frac{\partial}{\partial b} \frac{e \phi}{T_i n_0}$$
 and the $k_\perp$ in ω_{ci} and $\sqrt{b}$ cancel each other, leading to a finite non-adiabatic ion response even in the electron scale ($b \gg 1$) (Smolyakov et al., PRL 2002; Hirose and Elia, PoP 2002)




Inverse Aspect Ratio Scans at $\rho=0.8$ Show Consistent Trend with UM




The Transition wavenumber, $(k_\theta \rho_s)_{TR}$, at $\rho = 0.8$ Scales with ϵ Consistently with UM



Ion and Electron Temperature gradient Scans at $\rho = 0.8$ Show Consistent Trend with UM




ϵ and Temperature gradient Scans at $\rho = 0.6$ Show Consistent Trend with UM




KBM/ITG/TEM are likely the Operating Micro-instabilities at $\rho = 0.3$

- Distinct differences are seen in response to η_e and η_i variations by modes at $k_\theta \rho_s < 0.3$ and at $k_\theta \rho_s \gtrsim 0.3$.
- For $k_\theta \rho_s < 0.3$ modes: either reducing (increasing) η_e or η_i alone being stabilizing (destabilizing); completely stabilized with η_e and η_i both reduced by 70%; γ being more sensitive to η_i than to η_e ; ω being in the ion direction
 - KBM likely: modes likely being driven by total pressure, supported by odd parity in A_{ll} eigenmode structure (not shown) with much enhanced B fluctuations consistent with the higher β at $\rho = 0.3$ than at $\rho = 0.8$ (1.2% vs 0.05%)
- For $k_\theta \rho_s \gtrsim 0.3$ modes: γ enhanced with η_e and η_i reduced both by 70% or η_i reduced alone by 50% or η_e increased alone by 20%; γ reduced with η_i increased alone by 20% or η_e reduced alone by 50%; forming lower-k peak in γ with reduced η_i or η_e
 - Consistent with ITG/TEM hybrid
 - Reducing η_e or increasing η_i alone making modes more ITG-like with lower γ , forming a lower-k γ peaks, and ω in the ion direction
 - Reducing η_i alone leading to significantly enhanced γ and ω switching to the electron direction, consistent with TEM
 - Large reduction of both η_e and η_i leading to significantly enhanced γ and ω switching to the electron direction, clearly evidence of density gradient driven TEM
 - Averaged electron bouncing frequency ~ 4 (C_s/a), much higher than γ and ω at $k_\theta \rho_s \gtrsim 0.3$ and the detrapping frequency at $\rho = 0.3$.
- Wide scans in η_e and η_i represent large uncertainties in inferred profiles




Electron-scale Modes at $\rho = 0.8$ are very Sensitive to η_e

- γ of electron-scale modes at $k_\theta \rho_s > 7$ is very sensitive to η_e (through a/L_{Te})
 - Up to a factor of 3 increase in γ at $k_\theta \rho_s \approx 44$ with a 20% increase in η_e ; more than 80% decrease in γ in the $k_\theta \rho_s$ range of 23 to 78 with a 20% reduction in η_e
 - T_e gradient is the main free energy driving the unstable modes in the electron scale
 - Much weaker dependence on η_i , only significantly affected at $k_\theta \rho_s < 40$ with a 70% reduction in η_i
- ω of electron-scale modes at $k_\theta \rho_s > 7$ has very different response to η_e in the intermediate k range (between ρ_i^{-1} and ρ_e^{-1}) and in the high k range (ρ_e scale and beyond)
 - Intermediate k range modes having ω flipped to significantly positive values (not significantly affected) with a 20% increase (reduction) in η_e , while high-k modes having more (less) negative ω , consistent with the typical electrostatic ETG mode
 - Much weaker dependence on η_i , with significant changes in mode ω seen at $k_\theta \rho_s < 40$ with η_i reduced by 70%
- Increasing η_i by 20% does not significantly affect either γ or ω
- ω is a continuous function of $k_\theta \rho_s$ at $k_\theta \rho_s > 7$, indicating the same branch of solution of the dispersion relation in the electron scale.
- The intermediate range small- ω ETG-drive instability has quite different parametric dependence than the typical ETG mode and should be categorized as a different instability.



Identification of UM is Supported with a group of equilibria with different profile combinations

- Four plausible equilibrium combinations (most importantly T_e , T_i and n_e profiles) in another ST40 hot ion plasma, 10014, (similar to shot 9831) were used for GS2 linear gyrokinetic stability analysis at $\rho = 0.8$ where UM is clearly identified for shot 9831
 - To count for the 20% uncertainties in inferred profiles with different nominal value combinations
 - Having largest variations in T_i/T_e , electron-ion collision frequency across the four cases
- Clear UM-type ω sign switching is seen in the ion scale for cases 1 and 2
 - The existence of UM supported by ϵ and η_e/η_i scans
- Intermediate range small- ω instability and the typical ETG mode still exist
 - Supported by η_e and η_i scans
 - ω discontinuities showing the existence of another ETG-like instability forming a local γ peak in the intermediate scale



Summary

- Linear gyrokinetic stability analysis with GS2 code has been carried out for ST40 hot ion plasmas.
- Linear growth rates of both ion- and electron-scale ($k_\theta \rho_s \gtrsim 0.2$) modes decrease from the edge toward the core of the plasma ($\rho = 0.3\text{--}0.8$).
- Complete suppression of electron-scale modes ($k_\theta \rho_s \gtrsim 5$) is found at $\rho = 0.3$.
- Ubiquitous mode (UM), ETG and an unidentified low-real-frequency electron-scale instability are found at $\rho \gtrsim 0.6$, with KMB/ITG/TEM most likely at $\rho = 0.3$, through parametric scans, i.e., temperature gradients and aspect ratio.
- The identification of UM is supported with a group of equilibria with different profile combinations, particularly T_e , T_i and n_e , given the $\sim 20\%$ experimental uncertainty of inferred profiles.

