

Properties of anomalous ion heating in lower hybrid wave sustained plasmas

K. Iwasaki, A. Ejiri, N. Tsujii, K. Shinohara, O. Watanabe, S. Jang, Y. Peng, Y. Ko Y. Lin, Y. Shirasawa, T. Hidano, F. Adachi and Y. Tian

The University of Tokyo

Background and Objectives


- In the TST-2 spherical tokamak device, non-inductive start-up experiment by Lower-Hybrid waves (LHW) is conducted. $T_e > 10$ eV and $T_i \sim 10$ eV is observed.
- The LHW can couple with electrons, but not with ions due to the relation between particle velocities and wave phase velocity. Thus, the LHW can directly heat electrons, not ions.
- Because density is very low, the effect of collision with electrons is considered to be small.
- Power modulation experiment was conducted to investigate the response of ion temperature to LHW.

TST-2


Main parameter

- Major radius $R_0 = 0.36$ m
- Minor radius $a = 0.23$ m
- Toroidal field $B_t = 0.16$ T
- Plasma current $I_p < 27$ kA
- Duration $t < 100$ ms
- Electron density $n_e < 1 \times 10^{18} \text{ m}^{-3}$


Overview of TST-2



There are two capacitively coupled combine antenna to excite LHW (200.1 MHz).




Location of two LHW antenna projected on toroidal cross section




Top launched antenna

Outboard launched antenna



Picture of each sight on toroidal cross section



Visible spectroscopy


- Czerny-Turner spectrometer
- 16 CH PMT
- 0.2 nm/CH, Measurable: 200-500nm
- I-V convertor : < 0.01 ms
- 8 sights ($R_{\text{tan}}=180\text{-}540\text{mm}$) are available to measure to change them for each shot
- Impurity : C^{2+} (CIII, 464.74 nm)

Profiles of electrons and ions

Density : center peak


T_e : Hollow (10-50 [eV])

T_i : Flat or hollow (~ 10 [eV])




Analyzed Waveforms

Modulation period of LHW power is 6 ms.




Reconstruction of magnetic flux surface from magnetic probes calculated at 60 ms.




Method

Because emission from LHW plasma is weak, signal is analyzed by using a cycle averaged method. (3 times)
Each phase of modulation is named A to F like right figure.




Results

Time development after LH power is ON.



Time development after LH power is OFF.




Discussion

Solve 1-dimensional heat diffusion equation

$$\frac{\partial}{\partial t} \tilde{n} T_i = \underbrace{\frac{1}{\rho} \frac{\partial}{\partial \rho} \left(\rho \chi_{\text{neo}} \tilde{n} \frac{\partial T_i}{\partial \rho} \right)}_{\text{heat diffusion}} + \underbrace{P_{\text{coll}} + P_{\text{adj}}}_{\text{collision with electrons}}$$

χ_{neo} : Neoclassical heat diffusion coefficient
 P_{coll} : Collision between electrons and ions
 P_{adj} : Adjustable power which reproduce experiments




[i] $P_{\text{adj}} = 0$

Dotted blue line : initial T_i profile (roughly same as C on above figure.)
Blue line : Developed T_i (thin : 0.05ms after, thick : 1 ms after)
Black line : T_e profile

T_i gets about 2 eV (< 10 eV) in steady state.


Comparison of absolute value of heat flux (Calculated 0.05ms after initial state)



- Power of heat diffusion is much higher than power of collision with bulk electrons.
- Collision cannot sustain measured T_i .

- High energy electrons generated by LHW is considered to be dense on peripheral region, but collision with this fast electrons is further small.
- Other terms which can cool ions are not included. (e.g. charge-exchange) T_i could be lower than 2 eV.

[ii] $P_{\text{adj}} = 1.5$ [kW] ($0.48 \leq R \leq 0.52$)




- Flat profile ($\sim 10\text{eV}$) is reproduced when proper power on peripheral region is added.
- More power makes hollow profile. (E)
- Power in center region leads to center peak profile, cannot make hollow profile.

Ion heating seems to occur on peripheral region.


Parametric Decay Instability (PDI)

In TST-2, PDI is often observed. Ion cyclotron (Landau) damping of a wave generated by PDI is one candidate for ion heating.



Power of lower sideband increase with toroidal magnetic field.

Spectrum near 200 MHz $\rightarrow$



Conclusion and Future work

- Ion temperature responds to LHW modulation at peripheral region, does not respond near magnetic axis.
- Collision with electrons is very weak to explain the experimental results, other ion heating mechanism is needed on peripheral region.
- PDI is one candidate for ion heating and investigating the correlation between them is future work.

- Ion temperature is expected to increase with toroidal magnetic field.