



Observation of whistler wave instability driven by temperature anisotropy of energetic electrons on EXL-50 spherical torus

Mingyuan Wang^{1,2*}, Yuejiang Shi^{1,2}, Jiaqi Dong^{1,2}, Wei Chen³, Adi Liu⁴, G Zhang⁴, Shikui Cheng^{1,2,3}, Mingsheng Tan⁵, Songjian Li^{1,2}, Shaocong Song^{1,2}, Tiantian Sun^{1,2}, Hanyue Zhao^{1,2}, Bing Liu, Xianli Huang^{1,2}, Yumin Wang^{1,2}, Yingying Li^{1,2}, Xianming Song^{1,2}, Minsheng Liu^{1,2}, Baoshan Yuan^{1,2}, Y-K Martin Peng^{1,2}, and ENN team.

1 Hebei Key Laboratory of Compact Fusion, Langfang 065001, China
2 ENN Science and Technology Development Co., Ltd., Langfang 065001, China
3 Southwestern Institute of Physics, Chengdu 610041, China

4. Department of Plasma Physics and Fusion Engineering, University of Science and Technology of China, Anhui Hefei 230026, China
5 Institute of Energy, Hefei Comprehensive National Science Center, Hefei 230031, China



E-mail of corresponding author: wmyuan@mail.ustc.edu.cn

Abstract Whistler waves driven by temperature anisotropy energetic electrons are observed in electron cyclotron wave (ECW) steady state plasmas on XuanLong-50 (EXL-50) spherical torus. The whistler waves have multiple frequency bands which are proportional to the Alfvén velocity. Increasing of the bulk electron density or decreasing of the energetic electron temperature anisotropy reduces the intensity of the whistler wave and decreases the loss of energetic electrons. Moreover, the small disruption of plasma caused by whistler waves is also observed, probably as a result of whistler wave affecting the equilibrium profile. Furthermore, the interaction of whistler waves with low frequency acoustic waves has been observed. These provide a novel experimental way to study wave-particle interactions and whistler wave control of runaway electrons on future fusion devices.

1. XuanLong-50 spherical torus and resonance condition

- ✧ XuanLong-50 spherical torus (EXL-50) does not have central solenoid
- ✧ EXL-50 uses three sets of electron cyclotron wave (ECW) systems (28 GHz) to heat the plasma, start-up and ramp-up plasma current ($T_{\perp e} > T_{\parallel e}$).
- ✧ Energetic electrons contribute the main plasma current (**Bump on tail**)




Figure1. The schematic diagram of EXL-50




Figure2. Schematic diagram of electron velocity distribution function on EXL-50.

✧ Electron-wave coupling can be expressed as

$$\omega - k_{\parallel}v_{\parallel} - k_{\perp}v_d - l\omega_{ce}/\gamma = 0 \quad (1)$$




Figure3. Hard X-ray spectrum (left) and calculated resonance curves for different k and Bt (right)

2 High-frequency magnetic probe

- ✧ A high-frequency magnetic probe (HFP) is developed on EXL-50 to measure high-frequency magnetic field fluctuation
- ✧ **1-200 MHz** magnetic field fluctuation will be measured by the high-frequency magnetic diagnostic.
- ✧ Significant frequency chirp is observed




Figure4. Schematic of the hardware for HFP diagnostic (50 Ω).






Figure5. HFP has a high signal-to-noise ratio (left) and observed frequency chirp (right)

3. Whistler wave



4. Others



- ✧ Whistler waves may induce plasma disruption
- ✧ Observation of nonlinear interactions of energetic particles with whistler wave and sound wave

5. Conclusion

- ✧ Developed a high-frequency magnetic probe
- ✧ Whistler wave is driven by temperature anisotropy of energetic electrons in steady state ECW plasmas
- ✧ The frequency of the whistler wave is between 30-120 MHz, and is proportional to the Alfvén velocity
- ✧ The whistler wave intensity is verified to be correlated with the density and $\frac{T_{\perp e}}{T_{\parallel e}}$
- ✧ Whistler waves may induce plasma disruption
- ✧ Nonlinear interactions between energetic particles, whistler wave and sound wave

6. References

1. Gary S P, Wang J. J. Geophys. Res. Space Physics, 1996, 101(A5): 10749-10754.
2. Garner R C, et al. Phys. Rev. Lett., 1987, 59(16): 1821.
3. Fülöp T, Pokol G, Helander P, et al. Phys. Plasmas, 2006, 13(6): 062506.
4. Pong D A et al. Phys. Rev. Lett., 2018, 120(15): 155002.
5. H. Ikegami, S. Aihara, M. Hosokawa, and H. Aikawa, Nucl. Fusion 13, 351 (1973).
6. Y. Shi et al 2022 Nucl. Fusion 62 086047
7. M. Wang et al 2022 Plasma Phys. Control. Fusion 64 075006
8. Wang M, et al. arXiv preprint arXiv:2208.05323, 2022.
9. Wang M, et al. arXiv preprint arXiv:2108.09648, 2021.