

Global gyrokinetic study of shaping effects on electromagnetic modes at NSTX aspect ratio

**A. Y. Sharma, M. D. J. Cole, T. Görler, Y. Chen, D. R. Hatch,
B. J. Sturdevant, R. Hager, S. Ku, W. Guttenfelder, A. Mishchenko,
C. S. Chang**

- Introduction
- Use of profile-consistent magnetic equilibria
- Benchmarking of an approximation for the inclusion of compressional magnetic perturbation $\delta B_{||}$ effects
- Shaping study based on NSTX EP H-mode discharge #132588
- Conclusion

- NSTX discharge #132588 is an Enhanced Pedestal H-mode (EP H-mode) discharge that features:
 - An absence of edge localized modes (ELMs);
 - Significantly improved energy confinement;
 - High plasma β ;
 - High plasma shaping.
- We have performed a shaping study based on NSTX EP H-mode discharge #132588.
- Local gyrokinetic simulations of NSTX discharge #132588 have found that $\delta B_{||}$ effects are important, so we include an approximation for $\delta B_{||}$ effects.
- In order to properly take into account finite-pressure effects, we use profile-consistent magnetic equilibria when performing scans over plasma β .



- The X-point Gyrokinetic Code (XGC) is a total- f global electromagnetic (EM) gyrokinetic particle-in-cell (PIC) code.
- It can model magnetically confined plasmas for:
 - Tokamaks (whole plasma volume from the magnetic axis to the material wall);
 - Stellarators (core region).
- Total- f EM XGC is unique:
 - Includes realistic divertor and limiter geometry;
 - Allows for non-Maxwellian plasmas;
 - Computes neoclassical and turbulent transport together with self-consistent 3D background and perturbed fields;
 - Uses a nonlinear Fokker-Planck-Landau collision operator;
 - Includes neutral particle recycling, and heat and torque sources.
- For details, see R. Hager, S. Ku, A. Y. Sharma et al., “Electromagnetic total- f algorithm for gyrokinetic particle-in-cell simulations of boundary plasma in XGC”, Phys. Plasmas (submitted).



- In the work presented here:
 - Only the tokamak δf EM capability is used;
 - Collisions are not included;
 - The SOL is not included.
- For further details, see:
 - A. Y. Sharma *et al.*, “Global gyrokinetic study of shaping effects on electromagnetic modes at NSTX aspect ratio with ad hoc parallel magnetic perturbation effects”, Phys. Plasmas (2022).



- We take into account finite-pressure effects by using profile-consistent magnetic equilibria.
 - When scanning over β , the magnetic equilibrium is reconstructed at each value of β such that the magnetic equilibrium pressure is consistent with the input pressure profile.
- We have a convenient workflow for generating these profile-consistent magnetic equilibria for any:
 - Aspect ratio A ;
 - Elongation κ ;
 - Triangularity δ .
- The present study is limited to up-down symmetric magnetic equilibria.
 - An extension to up-down asymmetric magnetic equilibria should be relatively simple.



Workflow:



1. Compute Fourier coefficients of a boundary from:
 - a. Major radius R ;
 - b. Minor radius a ;
 - c. Elongation κ ;
 - d. Triangularity δ .
2. Provide VMEC with
 - a. Boundary Fourier coefficients;
 - b. Pressure profile;
 - c. q profile;
 - d. Enclosed magnetic flux.
3. Convert the VMEC output to G EQDSK format using the MConf tool (available at PPPL).
4. Convert from G EQDSK format to EQD format using a simple XGC tool.

Perturbed EM field model	Numerical implementation	Computational cost	Validity	Common usage
$\delta\phi, \delta A_{ }, \delta B_{ }$	Relatively complex	Relatively expensive	Always valid	Local simulations
$\delta\phi, \delta A_{ }, \delta B_{ }=0$	Relatively simple	Relatively cheap	Maybe invalid at: <ul style="list-style-type: none">• high β;• low aspect ratio.	Global simulations
$\delta\phi, \delta A_{ }$, approximate $\delta B_{ }$ effects	Relatively simple	Relatively cheap	Limited verification with local simulations	Local simulations



- To lowest order in β and $k_{\perp} \rho_i$, the inclusion of $\delta B_{||}$ cancels the pressure-dependent part of the ∇B -drift (Joiner, PoP, 2010).
- Thus, a lowest-order approximation for the inclusion of $\delta B_{||}$ effects is to remove the pressure-dependent part of the ∇B -drift.
- This approximation has previously been verified with local gyrokinetic codes.
- We have performed a benchmark of this approximation between the global gyrokinetic code XGC and the global versions of the gyrokinetic codes GEM and GENE.
- The case that is used is based on the Cyclone Base Case (CBC), except that the ratio of the ion gyroradius to tokamak minor radius ρ^* is approximately 1/50, compared to approximately 1/180 for the CBC, thus, we refer to this case as the Economical Cyclone Base Case (ECBC).
- A single toroidal mode number $n=6$ is considered, and real electron mass is used.

- Reasonably good agreement between the three codes.
- Stronger disagreement at 5 times the critical β .
 - This is perhaps unsurprising, and may just be a question of convergence.






- We have performed a shaping study using:
 - Profile-consistent magnetic equilibria;
 - Approximate compressional magnetic perturbation $\delta B_{||}$ effects;
 - Magnetic geometries that are based on NSTX EP H-mode discharge #132588;
 - Magnetic field strength and q profile from NSTX discharge #132588;
 - Plasma profiles that are based on the CBC;
 - A single toroidal mode number $n=20$;
 - Real electron mass.

It would also be interesting to consider other plasma profiles and mode numbers.



We consider the following geometries:

Geometry	Aspect ratio A	Elongation κ	Triangularity δ
Circular	1.4	1.0	0.0
Elongated	1.4	2.2	0.0
Elongated, triangular	1.4	2.2	0.55



- All of the geometries use an aspect ratio that is identical to that of NSTX discharge #132588.
- The “elongated” and “elongated, triangular” geometries use an elongation κ that is identical to that of NSTX discharge #132588.
- The “elongated, triangular” geometry uses a triangularity δ that is identical to that of NSTX discharge #132588.
- While NSTX discharge #132588 has some up-down asymmetry, all of these geometries exclude up-down asymmetry.




- Transitions from collisionless trapped electron modes (CTEMs) to collisionless microtearing modes (CMTMs) to KBMs are observed.
- CMTMs are shifted to higher β_e , and then suppressed by strong triangularity.
- Strong NSTX shaping makes KBM growth rates as weak as CTEM growth rates.




Mode structures for the electrostatic potential ϕ (left) and the parallel magnetic potential $A_{||}$ (right) for the CMTM observed with the “circular” geometry case at $\beta_e = 10\%$.




Mode structures along a field line for the electrostatic potential ϕ (left) and the parallel magnetic potential $A_{||}$ (right) showing tearing parity for the CMTM observed with the “circular” geometry case at $\beta_e=10\%$.




- Transitions from collisionless trapped electron modes (CTEMs) to collisionless microtearing modes (CMTMs) to KBMs are observed.
- CMTMs are shifted to higher β , and then suppressed by strong triangularity.
- Strong NSTX shaping makes KBM growth rates as weak as CTEM growth rates.



Mode structures for the electrostatic potential ϕ (left) and the parallel magnetic potential $A_{||}$ (right) for the CTEM observed with the “elongated” geometry case at $\beta_e = 15\%$.



- Transitions from collisionless trapped electron modes (CTEMs) to collisionless microtearing modes (CMTMs) to KBMs are observed.
- CMTMs are shifted to higher β , and then suppressed by strong triangularity.
- Strong NSTX shaping makes KBM growth rates as weak as CTEM growth rates.



Mode structures for the electrostatic potential ϕ (left) and the parallel magnetic potential $A_{||}$ (right) for the KBM observed with the “elongated, triangular” geometry case at $\beta_e = 30\%$.

- A workflow has been developed for performing scans over plasma β using profile-consistent magnetic equilibria with the global gyrokinetic PIC code XGC.
- An approximation for the inclusion of compressional magnetic perturbations $\delta B_{||}$ was benchmarked with the global gyrokinetic code XGC and the global versions of the gyrokinetic codes GEM and GENE.
- The effects of high plasma shaping at NSTX aspect ratio are that there is significant stabilization of CMTMs and KBMs.
 - KBM growth rates are reduced to that of the TEMs.